
## **Polar Rock Repository**





- •Established in late 2003 to preserve polar rock and dredge samples for future research, education & exhibit use
- •The PRR provides full and open access to samples and metadata via the website.







Samples are weighed, photographed, measured for susceptibility & put into bags with QR code labels

•••• AT&T **?** 

8:19 AM

**1** 99% €






Period

http://research.bpcrc.osu.ed...












Multimedia						
Primary Field #	97-13-5					
IGSN #	IGSN.PRR036239					
Kind of Object	Rock Sample					
Rock Type	Igneous					
Rock Name	Breccia Breccia					
Sample Description	Friable, clasts to 10cm, matrix includes quartz, ash (basaltic).					
Weight	2157 g					
Susceptibility	55					
Notes	Consists entirely of coarse breccia, broken only by a dolerite intrusion halfway up south ridge (which is where section measured). Vertical interval measured parallel to strike and assuming a 20 degree dip measured on similar breccias on next outcrop to NW).					
Additional Notes at PRR	Yes					
Continent	Antarctica					
Bogien	Couthorn Victoria Land					

Range/Island Group	A CONTROL OF THE CONT
Mountain/Island Name	Ricker Hills
Precise Location	Most south of low outcrops between Ricker Hills and Thomas Rock (and outcrop closest to camp)
Coordinates	-75.703, 158.784 (75° 42.2' S, 158° 47' E)
Positioning Method	Notes

Epoch	Early							
Stratigraphy	Group	Formation	Member	Section Name	Unit Thickness	Sample Height	Distance from Intrusion	
		Mawson Formation				24 m		

Primary Field Logistics	LC130 Hercules Aircraft				
Data Sources	Field Notebook				
Field Year Collected	1997				
Collector	Elliot, David H.				



New collections: Eltanin Cruises Dredge samples, Shackleton Glacier (Beacon), Scott & Reedy glaciers

- Online access to >42,000 rock samples
- Samples freely available for research and education